




Potency of Poxviral Based Vectors for Malaria and TB vaccines

Anne Moore

Adrian Hill

**Wellcome Trust Centre for Human Genetics
Oxford University**


Malaria Vaccines in Clinical Development



Pre-erythrocytic Strategies


- Prevent sporozoite invasion of liver cell:
Antibody
- Prevent sporozoite surviving, replicating in and progressing from liver cell:
T cell

Recombinant Malaria Vaccines




CSO construct: codon optimised *PfCS*, with C-term Pb9 (PbCSP epitope)

Recombinant Poxviruses



MVA

- ❖ Highly attenuated vaccinia virus strain
- ❖ Deletions in host range genes and cytokine genes
- ❖ Replicates in chick embryo fibroblasts but not in mammalian cells
- ❖ Excellent safety record - over 120,000 people immunised with MVA




Fowlpox9


- ❖ An avipoxvirus, like canarypox
- ❖ Attenuated by Anton Mayr (Munich)
- ❖ Characterised molecularly by the Institute of Animal Health, Compton, UK (Mike Skinner)
- ❖ Shown in Oxford to be more immunogenic for CD8+ T cell induction than the standard fowlpox strain

Malaria Vaccines in Mice

Pb9-specific CD8⁺ T-cells in Spleen, Liver and Blood



Protective efficacy of prime-boost immunisation regimes




Prime / Boost	Animals Protected	Efficacy (%)
FP / MVA	27 / 40	67.5
MVA / FP	15 / 40	37.5
FP / FP	3 / 19	15.7
MVA / MVA	3 / 20	15.0

2000 sporozoites


Oxford University Malaria Vaccine Trials

		<u>Vaccinees</u>	<u>Challengees</u>	<u>Year</u>
VAC 01	DNA im vs gene gun +/- MVA boost id	12		1999
VAC 02	MVA id	6		
VAC 03	DDDM, DMM, DDD, MMM	15		2000
VAC 04	challenge study for 03		17	
VAC 05	DDMM vs GGMM	9		
VAC 06	<i>Gambia: DDMM vs MMM</i>	18		
VAC 07	challenge study for 05		14	
VAC 08	MVA boosting of sporozoite challengees	4		
VAC 09	MVA higher dosages	6		2001
VAC 10	DDMM higher dose, longer interval	9		
VAC 11	<i>Gambia: 1-5 year old children MVA</i>	20		
VAC 12	FP9 phase I and high dose MVA boost	12		
VAC 13	challenge studies for 10 and 12		14 + 10	
VAC 14	<i>Gambia: FFM, DDM, DDDDM high dose</i>	30		
VAC 15	FFM vs MMM high dose	17		2002
VAC 16	challenge study for 15		22	
VAC 17	FM, MF, DDFM, DDMF	17	23	
VAC 18	RTS,S MVA-CS, MVA-CS RTS,S	24	17	
VAC 19	<i>Kilifi, Kenya FM, FFM</i>	30		
VAC 20	<i>Gambia DDM vs rabies</i>	328		

Immunogenicity in Man: *Ex Vivo* Elispot Responses; Oxford-based trials




Malaria Vaccine Efficacy in Oxford Sporozoite Challenge Studies




All vaccines express ME-TRAP

*Bejon et al. *J. Infect. Dis.* (2004)

Cultured ELISPOT Responses




Correlation with Delay to Parasitemia



Spearman $r = 0.499$; $p = 0.015$

TB trials

Median Elispot summed peptide pool responses:
BCG v MVA85A v BCG-MVA85A



Conclusions from Vaccine Studies

- Heterologous prime-boost regimes that work in mice also work in humans. Murine model can predict vaccine efficacy.
- Correlates of immunity? Malaria: Memory T cell responses (cultured elispot at day of challenge) strongly correlate with delay to parasitemia. TB: CD4+ T cell responses?

Phase I & IIa/b Clinical Vaccine Production & Release

- Oxford:
- Candidate Recombinant Poxvirus Vaccine:
 - Confirm insert sequence, identity & purity PCR (absence of WT virus)
 - Confirm immunogenicity: Potency assay in mice (IFN- γ elispot)

- IDT,
Germany
- GMP manufacture of clinical lot:
 - QA/QC: Purity,
Viral Titre, live infectious virus
Lack of Extraneous agents, Endotoxin, Mycoplasma
Lack of Replication Competence

- HLS
- Toxicology
 - Persistence & distribution

- Oxford:
- QA/QC:
Purity, Insert Sequence,
Viral Titre,
Immunogenicity; Potency assay

Batch released for clinical use.

Potency Assay: IFN- γ ELISPOT Assay


4 mice immunized,
Antigen-specific
response >
15sfc/million, no more
than 1 non-responder

Antigens tested

P. berghei MHC class I
Peptide; Pb9 SYIPSAEKI

OR

M.tb85A
MHC class I & II epitope



Phase III & Beyond...

- Dose titration potency assay; increase sensitivity; detect subtle differences between lots
- 3Rs; should move away from routine potency testing in animals
- Alternative: does *in vitro* mRNA or protein expression correlate with immunogenicity *in vivo*?
- Tests to quantify the protein expression to be implemented

Future Directions in Production & Validation of Recombinant Virus Vector-based Vaccines

- New antigens or combinations of expressed antigens; Polyprotein construct; new challenges with larger inserts.
- New viral vectors; Adenoviruses
- New GMP facilities; Oxford GMP facility to produce clinical lots of recombinant ADV-METRAP & poxvirus-based vaccines.

Acknowledgements

Oxford Pre-Clinical

Sarah Gilbert
Claire Hutchings
Simon Draper
Eric Prieur
Richard Anderson
Joerg Schneider
Simone Korten
Kate Watkins

NMRC

Judy Epstein
Dan Carucci

WRAIR

Jim Cummings
Kent Kester
Jack Williams
Gray Heppner

MRC, The Gambia

Vasee Moorthy
Kalifa Bojang
Paul Milligan
Margaret Pinder
Tunde Imoukhuede

LSHTM

Brian Greenwood

Kilifi, Kenya

Philip Bejon
Kevin Marsh

GSK

Joe Cohen
Amanda Leach
Rip Ballou

Oxford Clinical Trials

Susie Dunachie
Michael Walther
Sam McConkey
Daniel Webster
Jenni Vuola
Sheila Keating
Tamara Berthoud
Laura Andrews
Ian Poulton
Rikke Esbensen
David Warrell
Frances Sanderson

Imperial College

Bob Sinden
Geoff Butcher

Oxford TB

Helen McShane
Ansar Pathan
Helen Fletcher

MVI at PATH, The Wellcome Trust